

AAS DPS 2008, Ithaca, NY, October 11–15

---

# Resonant Asteroid Families

## a Wealthy Source of Information on Planetary Migration

---

M. Brož<sup>1</sup>, D. Vokrouhlický<sup>1</sup>

<sup>1</sup> Charles University, Prague


---

### Table of contents:


- (PART 1) asteroid families in the J3/2 resonance
- (PART 2) resonant Yarkovsky effect, age determination
- (PART 3) are there any Trojan families?

# Update of resonant populations (Aug 2007):

**270** Zhongguos, Griquas and unstable    **1200** Hildas    **3** Thule




# J3/2 — pseudo-proper elements:




- a distinct collisional **Schubart family** (Schubart 1991);  
 $v_{\text{cutoff}} \simeq 60$  m/s,  $\sim 100$  km parent body, LF/PB  $\simeq 0.25$ ,  
 C/X-type members, SFD steeper than J3/2.
- **Hilda family** at higher inclinations;  $\sim 200$  km PB

# Simulated impact in J3/2:



- a disruption of a 100 km PB; isotropic velocity field with  $v_{\text{mean}} = 50 \text{ m/s}$ ,  $f = 0$ ,  $\omega + f = 180^\circ$
- ‘mapping’ of the osculating elements into pseudo-proper


# Impact in J3/2 — evolution over 4 Gyr:



- $N$ -body simulation, SWIFT integrator, 4 planets only
- the shape of the swarm is well **preserved** for 4 Gyr
- problem: observed **Schubart family** has larger spread in  $e$




# Impact in J3/2 — Yarkovsky effect:




- the Yarkovsky induces the **drift in eccentricity**  
→ different mechanism than for Main Belt families!

# Yarkovsky **drift** in $e$ (RTBP):



- object is locked in the libration centre,  $a$  is almost fixed
- adiabatic invariant  $N = \sqrt{a} \left( \frac{p+1}{p} - \sqrt{1-e^2} \cos i \right)$  evolves linearly


# Schubart and Hilda families — ages:



- K-S test of  $e$ -distributions for Schubart  $\Rightarrow$  1–2.5 Gyr old
- $\gtrsim 4$  Gyr for Hilda (LHB origin?)




# Hildas vs 2:1 Jupiter–Saturn resonance:




- **J3/2 population is strongly unstable** during 2:1 resonance crossing (Brož & Vokrouhlický 2008)
- all families have to be younger than the time of 2:1 (collisional probabilities might be higher early after 2:1)

# Hilda family — ( $e$ , $H$ ) plot:




- ‘ears’ like in ( $a$ ,  $H$ ) for Eos, Erigone, Massalia, Merxia, ...
- **YORP effect** changes spins  $\Rightarrow$  more precise age?

# Trojans — how many families?:




- $\sim 1000$  bodies in each  $L_4$  and  $L_5$  cloud
- definitely not many prominent clusters...

# Trojans — only 1 family in L4?:



- $N(v_{\text{cutoff}})$  plots compared with (random) background
- **Eurybates family** is the most robust case; it has only C-type members, no D-types (Roig *et al.* 2008)


# Eurybates family — SFD:



- only Eurybates has steeper SFD than the background  
 $\Rightarrow$  consistent with collisional origin
- 1 collisional family among Trojans consistent with J3/2  
 $\Leftarrow$  mean collisional probabilities are comparable:  
 $16$  and  $11 \times 10^{-18} \text{ km}^{-2} \text{ yr}^{-1}$  (Dahlgren 1998)




# Eurybates family — no Yarkovsky effect:



- evolution with/without Yarkovsky effect is the same (even for strengthened thermal forces)
- different dynamics in 0th order resonance  $\Rightarrow$  no systematic drift in  $e$ , no precise age



# Eurybates family — chaotic diffusion:



- present shape was attained due to the chaotic diffusion
- age might be 1–4 Gyr (‘filament’ must disappear)


## Conclusions:

- resonant Yarkovsky effect operates
- two collisional families in the J3/2 resonance  
(probably old, shallower SFD — different from MB)
- one collisional family among Trojans

## Future work:


- $(e, H)$  method  $\Rightarrow$  more precise ages of resonant families
- stability of resonant families during planetary migration
- repopulation efficiency from the planetesimal disc

# Schubart family — colours:




- SLOAN colours of J3/2 bimodal (C/X- and D-types)
- Schubart cluster only C/X-type  $\Rightarrow$  collisional origin

# Schubart family — SFD:




- Schubart family significantly **steeper** than the rest of J3/2  
 $\Rightarrow$  collisional origin (but not as steep as MB families)

# A comparison with Merxia family ( $a, H$ ):



# Hilda — expected $\Delta a$ vs observed $\Delta e$ :



- linear dependence  $\Rightarrow$  the same code as for  $(a, H)$